


المادة: الرياضيات الشهادة: المتوسطة نموذج رقم -3- المدة : ساعتان	الهيئة الأكاديمية المشتركة قسم : الرياضيات	 المركز التربوي للبحوث والإنماء
---	---	---

نموذج مسابقة (يراعي تعليق الدروس والتوصيف المعدل للعام الدراسي ٢٠١٦-٢٠١٧ وحتى صدور المناهج المطورة)

ارشادات عامة: - يسمح باستعمال آلة حاسبة غير قابلة للبرمجة او اختزان المعلومات او رسم البيانات.  
 - يستطيع المرشح الإجابة بالترتيب الذي يناسبه دون الالتزام بترتيب المسائل الوارد في المسابقة.

### I- (3 points)

Answer « True » (T) or « False » (F) and justify your answer.

- 1)  $(-2x - 2)^2 = 4(x + 1)^2$ .
- 2) The solutions of the equation  $x^2 + 10 = 0$  are  $\sqrt{10}$  and  $-\sqrt{10}$ .
- 3) If  $x$  is an acute angle and  $\sin x = \frac{1}{3}$ , then  $\cos x = \frac{2}{3}$ .
- 4) The equation  $(x + 3)^2 = 0$  has no solution.
- 5) If  $x$  is a number greater than 3, then  $(x^2 + 1)(2x - 5)$  is positive.

### II- (2 points)

The questions 1) et 2) are independent. Show all the steps of your work.

1) Given  $A = \frac{1}{\sqrt{7}+1} + \frac{1}{\sqrt{7}-1}$  and  $B = \frac{7}{3\sqrt{7}}$ .

Compare A and B.


2) a) Verify that:  $\frac{4\sqrt{2}+2}{4+\sqrt{2}} = \sqrt{2}$ .

b) Use the previous equality to prove that  $\frac{(\sqrt{32}+2)^2}{(\sqrt{36}-10-\sqrt{2})^2}$  is a natural number.

### III- (4 points)

ABCD is a rectangle such that  $AB = 4$  m and  $AD = 3$  m.

M is a point on [AD]. The parallel through D to (BM) intersects with [BC] at N. Let  $AM = x$ .



#### Part A

- 1) Prove that:
  - a)  $x$  is less than 3.
  - b) DMBN is a parallelogram.
  - c)  $NC = x$ .
- 2) Prove that the area of the square with side DM is  $(3 - x)^2$ .
- 3) Prove that the area of the parallelogram DMBN is equal to  $12 - 4x$ .

#### Part B

- 1) Factorize  $S' - S$ .
- 2) Can you find  $x$  so that the two areas are equal?
- 3) a) Solve the equation  $(x+1)(3-x) = 3$ .  
 b) Give a geometric interpretation to the result.

#### IV- (2 points)

The sum of two numbers is 47. When we divide one of the numbers by 2 and the other by 3, the sum becomes 17.5.

1) Which one of these 3 systems is related to the given?

$$\begin{cases} x + y = 47 \\ 3x + 2y = 17.5 \end{cases} \quad \begin{cases} y = 47 - x \\ 3x + 2y = 105 \end{cases} \quad \begin{cases} x + y = 47 \\ \frac{x + y}{2} + \frac{y}{3} = 17.5 \end{cases}$$

2) Find the two numbers.

#### V- (4 points)

In an orthonormal system of axes  $x'Ox$  and  $y'Oy$ , consider the line (D) with equation  $y = 2x - 1$ , and the points B(2 ;3) and C(3 ;1).

1) Draw the line (D) and plot the points B and C.

2) Does the line (D) pass through the points B and C? Justify.

3) Let (D') be the line with equation  $y = -\frac{1}{2}x + \frac{5}{2}$ .

a) Prove that (D') passes through C and is perpendicular to (D).

b) (D') and (D) intersect at S. Determine the coordinates of the point S.


4) Determine the coordinates of the point I, centre of the circle circumscribed about the triangle BSC, and determine the length of its radius.

5) Determine the coordinates of point A such that BSCA is a parallelogram. Prove that A is a point on the circle circumscribed about BSC.

#### VI- (5 points)

In the adjacent figure :

- a semicircle with diameter [AB] and center O;
- $AB = 2R$ ;
- E is midpoint of [OB];
- (GE) perpendicular bisector of [OB] (G is a point on the semicircle);
- K is a point on segment [EG]. The line (BK) and the semicircle intersect at M.



1) Draw a figure, to be completed in the remaining parts of the problem.

2) a) Prove that the triangle OBG is equilateral.

b) Calculate GE in terms of R.

c) Calculate the angle GMB.

3) Prove that the triangles BEK and BMA are similar.  
Deduce that  $BK \times BM = R^2$ .


4) The perpendicular through E to (AM) intersects (AM) at N.

Calculate the ratio  $\frac{MN}{AM}$

5) In this part, suppose that k is the centroid of the triangle GOB.

a) Calculate EN and MN in terms of R.

b) Find R so that the perimeter of the quadrilateral BMNE is equal to  $7\sqrt{3} + 3$

المادة: الرياضيات الشهادة: المتوسطة نموذج رقم -3- المدة : ساعتان	الهيئة الأكاديمية المشتركة قسم : الرياضيات	 المركز التربوي للبحوث والإنماء
---	---	---

أسس التصحيح (تراعي تعليق الدروس والتوصيف المعدل للعام الدراسي ٢٠١٦-٢٠١٧ وحتى صدور المناهج المطورة)

### Answer keys

<b>I.</b>	1)	True Expand both expressions or: $(-2x - 2)^2 = [-2(x + 1)]^2 = 4(x + 1)^2$	<b>0.5</b>
	2)	False The equation has no solution : a square can't be negative .	<b>0.5</b>
	3)	False $(\cos x)^2 = 1 - (\sin x)^2 = \frac{8}{9}$ ; $\cos x = \frac{2\sqrt{2}}{3}$	<b>0.75</b>
	4)	False only -3 is the solution.	<b>0.5</b>
	5)	True because : $x^2 + 1 > 0$ for all $x$ ; and $2x - 5 > 0$ and $x > 2,5$ . The product of two positive numbers is positive.	<b>0.75</b>
<b>II.</b>	1)	$A = \frac{2\sqrt{7}}{6} = \frac{\sqrt{7}}{3} = \frac{\sqrt{7} \times \sqrt{7}}{3\sqrt{7}} = \frac{7}{3\sqrt{7}}$ , hence $A = B$	<b>0.75</b>
	2-a	we can show that: $\sqrt{2} \times (4 + \sqrt{2}) = 4\sqrt{2} + 2$	<b>0.5</b>
	2-b	$\frac{(\sqrt{32} + 2)^2}{(\sqrt{36} - 10 - \sqrt{2})^2} = \frac{(4\sqrt{2} + 2)^2}{(6 - 10 - \sqrt{2})^2} = \frac{(4\sqrt{2} + 2)^2}{(-4 - \sqrt{2})^2} = \frac{(4\sqrt{2} + 2)^2}{(4 + \sqrt{2})^2} = \left(\frac{4\sqrt{2} + 2}{4 + \sqrt{2}}\right)^2 = (\sqrt{2})^2 = 2$ , and 2 is a natural number.	<b>0.75</b>
<b>III.</b>	A.1-a	AD = 3 and $x$ is positive, hence $x$ is between 0 and 3.	<b>0.25</b>
	A.1-b	DMBN is a parallelogram since opposite sides are parallel.	<b>0.5</b>
	A.1-c	AD = BC , since ABCD is a rectangle ; and DM = NB, since DMBN parallelogram. Therefore AD - DM = BC - NB, and AM = NC = $x$	<b>0.5</b>
	2)	DM = 3 - $x$ , then area of the square = $(3 - x)^2$ .	<b>0.25</b>
	3)	Different ways: Area (DMBN) = Area (ABCD) - 2 x Area (AMB), because the triangles AMB and DCN are congruent. Area (DMBN) = 12 - 4 $x$	<b>0.5</b>
B-1	$(12 - 4x) - (3 - x)^2 = (3 - x)(x + 1)$ .	<b>0.5</b>	

	<b>B-2</b>	$x=3$ or $x = -1$ , both are rejected.	<b>0.5</b>
	<b>B-3-a</b>	$(x+1)(3-x) = 3$ , then $x=2$ or $x=0$ .	<b>0.5</b>
	<b>B-3-b</b>	The area of parallelogram is 3 more than the area of square.	<b>0.5</b>
<b>IV</b>	1)	$\begin{cases} x + y = 47 \\ \frac{x}{2} + \frac{y}{3} = 17.5 \end{cases}$ same as $\begin{cases} y = 47 - x \\ 3x + 2y = 105 \end{cases}$	<b>1</b>
	2)	The two numbers are 11 and 36	<b>1</b>
<b>V</b>	1)	Figure.	<b>0.5 + 0.25</b>
	2)	For $x = 2$ , $2x - 1 = 3$ ; hence B is on (D). For $x = 3$ , $2x - 1 = 5$ ; hence C is not on (D).	<b>0.25</b> <b>0.25</b>
	3) a	For $x = 3$ , $-\frac{1}{2}x + \frac{5}{2} = 1$ ; hence C is on (D').	<b>0.25</b>
	3) b	$\begin{cases} y = 2x - 1 \\ y = -\frac{1}{2}x + \frac{5}{2} \end{cases}$ therefore $2x - 1 = -\frac{1}{2}x + \frac{5}{2}$ ; $x = \frac{7}{5}$ et $y = \frac{9}{5}$ and $S(\frac{7}{5}; \frac{9}{5})$	<b>0.25</b> <b>0.5</b>
	4)	I midpoint of [BC]. Therefore $I(\frac{5}{2}; 2)$ $R = IB = \frac{\sqrt{5}}{2}$	<b>0.25</b> <b>0.5</b>
	5)	$\vec{BA} = \vec{SC}$ , therefore : $x - 2 = 3 - \frac{7}{5}$ and $y - 3 = 1 - \frac{9}{5}$ $x = \frac{18}{5}$ and $y = \frac{11}{5}$ BSCA rectangle, so BSC is a right triangle at A; and A is on the circle circumscribed about triangle BSC	<b>0.5</b> <b>0.5</b>
<b>VI</b>	1)		<b>0.5</b>
	2)a-	OG = BG (perpendicular bisector); OG = OB (radii) Therefore OG = BG = OB, and OBG equilateral.	<b>0.75</b>
	2) b-	$GE = \frac{R\sqrt{3}}{2}$ .	
	2) c-	$\widehat{GMB} = \frac{\widehat{GOB}}{2} = 30$ degree.	
	3)	BMA and BEK are right triangles and they have a common angle $\hat{B}$ . $\frac{BA}{BK} = \frac{AM}{KE} = \frac{MB}{EB};$ therefore $BK \times BM = BA \times EB = 2R \times \frac{R}{2} = R^2$	<b>0.75</b> <b>0.5 + 0.5</b>
4)	$\frac{MN}{AM} = \frac{BE}{BA} = \frac{1}{4}$ .	<b>0.75</b>	
5-a	$AM = \frac{AB}{2} = R$ and $EN = \frac{3R\sqrt{3}}{4}$ .	<b>0.5</b>	

	<p><b>5-b</b></p> $BE = \frac{R}{2}$ $MN = \frac{R}{4} \text{ and } MB = R\sqrt{3}$ $\frac{3R\sqrt{3}}{4} + \frac{2R}{4} + \frac{R}{4} + \frac{4R\sqrt{3}}{4} = 7\sqrt{3} + 3$ $R = 4.$	<p><b>0.5 +</b> <b>0.25</b></p>
--	---	-------------------------------------